$(2+1)-$dimensional supergravity invariant under AdS–Lorentz superalgebra.

O. Fierro[1], F. Izaurieta[1], P. Salgado[1], O. Valdivia[1,2]

[1]*Departamento de Física, Universidad de Concepción,*

Casilla 160-C, Concepción, Chile. and

[2]*Department of Mathematics, Heriot-Watt University,*

Colin Maclaurin Building, Riccarton, Edinburgh EH14 4AS, U.K.

Maxwell Institute for Mathematical Sciences, Edinburgh, U.K.

arXiv:1401.3697v2 [hep-th] 7 Mar 2014

Abstract

It is shown that the AdS–Lorentz superalgebra proposed in Ref. [1] can be obtained from the orthosymplectic algebra $OSp(N,4)$ (Anti-de-Sitter Superalgebra) following the procedure used in Ref. [2]. The S-expansion mechanism is used to obtain the (i) Casimir operators for the semi-simple extended Poincare superalgebra and (ii) the corresponding invariant tensors which allows us the construction of a $(2+1)$-dimensional Chern–Simons Supergravity Lagrangian.

CONTENTS

I. **Introduction** 3

II. **The AdS–Lorentz superalgebra** 4

III. **The S-Expansion of the AdS superalgebra** 6

A. **Semigroup S_{S3}** 7

B. **Semigroup S_{S2}** 9

C. **AdS–Lorentz Casimir operators** 12

IV. **AdS–Lorentz Chern-Simons supergravity in $D = 3$** 15

A. **The Invariant Tensor** 15

B. **AdS–Lorentz invariant supergravity action** 16

V. **Comments** 17

References 17

I. INTRODUCTION

The Maxwell algebra was introduced in the early seventies [3],[4] but it is only recently that has attracted more attention [5], [6], [7], [8], [9]. This algebra describes the symmetries of a particle moving in a background with a constant electromagnetic field.

The renewed interest in the Maxwell symmetry is due to its supersymmetrization leads to a new $N = 1$, $D = 4$ superalgebra, containing the Poincare superalgebra as its subalgebra [6]. Even more interestingly, in Ref. [9] it has been argued that the gauged Maxwell algebra generalizes the cosmological term in Einstein gravity. The supersymmetric version, on the other hand, seems to be specially appealing and with a beautiful interpretation: It describes the supersymmetries of $N = 1$, $D = 4$ superspace in the presence of a constant abelian supersymmetric field strength background [6]. The anti de–Sitter (AdS) counterpart of the Maxwell algebra namely the AdS–Maxwell algebra, and its supersymmetric extension, have also been considered in [10] and [11] respectively. The Maxwell algebra and its AdS–Maxwell analogue are both non semisimple, and therefore they have a degenerate Cartan–Killing metric. However, in ref. [1] a semisimple tensor extension of the Poincare algebra, which we call AdS–Lorentz algebra, is proposed. Following this reference, it is also shown that, in $D = 4$ dimensions, the AdS–Lorentz algebra admits $N = 1$ supersymmetric extension. The AdS–Lorentz algebra has the peculiarity that it can be obtained as an S–expansion mechanism [12], [13], [14], [15], [16] of the AdS algebra $SO(D-1,2)$. S-expansions are systematic ways to enhance symmetries and they are particularly interesting because they provide of the invariant tensors associated to the expanded algebra. This is a key ingredient in the construction of Chern–Simons theories.

In this work, three goals are achieved: First of all it is shown that the AdS–Lorentz superalgebra can be obtained as an S–expansion mechanism of the orthosymplectic algebra $OSp(N,4)$. Secondly, the invariant tensors of the AdS–Lorentz superalgebra are computed and the Casimir operators given in [1] are re–obtained. Finally, we construct the associated $D = 2+1$ dimensional Chern-Simons supergravity action. The discussion throughout this paper is organized as follows: In Section II we introduce the AdS–Lorentz superalgebra. In Section III we briefly review the main aspect of the S–expansion procedure and it is shown that the AdS–Lorentz superalgebra $SO(2,1) \oplus OSp(2|1) \otimes Sp(2)$ can be obtained from the $N = 1$ anti-de-Sitter super algebra $OSp(2|1) \otimes Sp(2)$ via the S–expansion procedure. In

section IV, we compute the Chern–Simons supergravity action in $(2+1)-$dimensions. A brief comment in section V concludes this paper.

II. THE ADS–LORENTZ SUPERALGEBRA

In $D = 4$ dimensions, the AdS–Lorentz algebra admits the following supersymmetric generalization [1],

$$[J_{ab}, J_{cd}] = \eta_{ad}J_{bc} + \eta_{bc}J_{ad} - \eta_{ac}J_{bd} - \eta_{bd}J_{ac} \tag{1}$$

$$[J_{ab}, P_c] = \eta_{bc}P_a - \eta_{ac}P_b \tag{2}$$

$$[P_a, P_b] = cZ_{ab} \tag{3}$$

$$[J_{ab}, Z_{cd}] = \eta_{ad}Z_{bc} + \eta_{bc}Z_{ad} - \eta_{ac}Z_{bd} - \eta_{bd}Z_{ac} \tag{4}$$

$$[Z_{ab}, P_c] = \frac{4a^2}{c}\left(\eta_{bc}P_a - \eta_{ac}P_b\right) \tag{5}$$

$$[Z_{ab}, Z_{cd}] = \frac{4a^2}{c}\left[\eta_{ad}Z_{bc} + \eta_{bc}Z_{ad} - \eta_{ac}Z_{bd} - \eta_{bd}Z_{ac}\right] \tag{6}$$

$$\{Q_\alpha, Q_\beta\} = -d\left[\frac{2a}{c}\left(\gamma^a C\right)_{\alpha\beta}P_a + \left(\sigma^{ab}C\right)_{\alpha\beta}Z_{ab}\right] \tag{7}$$

$$[J_{ab}, Q_\alpha] = -\left(\sigma_{ab}Q\right)_\alpha \tag{8}$$

$$[P_a, Q_\alpha] = a\left(\gamma_a Q\right)_\alpha \tag{9}$$

$$[Z_{ab}, Q_\alpha] = -\frac{4a^2}{c}\left(\sigma_{ab}Q\right)_\alpha \ , \tag{10}$$

Here Q_α are the super translation generators and Z_{ab} is the second rank antisymmetric tensor which characterizes the extension. Moreover a, c and d are arbitrary real constants and $\sigma_{ab} = \frac{1}{4}[\gamma_a, \gamma_b]$, where the γ_a are Dirac matrices. In the limit $a \to 0$, $\mathcal{N} = 1$, the standard Maxwell superalgebra is recovered.

The superalgebra (1)-(10) has been addressed in different ways in literature. Throughout this work, we call it AdS–Lorentz superalgebra.

In order to obtain the AdS–Lorentz superalgebra in $D = 3$ dimensions, a possible way to do so is to introduce arbitrary parameters in the (anti)-commutation relations similar to

the ones for $D = 4$ dimensions,

$$\{Q_\alpha, Q_\beta\} = -d\left[\mu\frac{2a}{c}\left(\gamma^a C\right)_{\alpha\beta} P_a + \nu\left(\sigma^{ab}C\right)_{\alpha\beta} Z_{ab}\right] \tag{11}$$

$$[J_{ab}, Q_\alpha] = \chi\left(\sigma_{ab}Q\right)_\alpha \tag{12}$$

$$[P_a, Q_\alpha] = \tau a\left(\gamma_a Q\right)_\alpha \tag{13}$$

$$[Z_{ab}, Q_\alpha] = \zeta\frac{4a^2}{c}\left(\sigma_{ab}Q\right)_\alpha . \tag{14}$$

Here μ, ν, χ, τ, ζ are arbitrary constants and we choose a representation for the Clifford algebra in three dimensions in terms of gamma matrices:

$$\begin{aligned}
&\gamma_0 = i\sigma_y = -C \ , \quad \gamma_1 = \sigma_x \ , \quad \gamma_2 = \sigma_z \\
&[\gamma_a, \gamma_b] = -2\epsilon_{abc}\gamma^c \ , \quad \{\gamma_a, \gamma_b\} = 2\eta_{ab} \ , \\
&\text{with} \quad \eta_{00} = -\eta_{11} = -\eta_{22} = -1 \\
&\text{and} \quad \gamma_a^T = -C^{-1}\gamma_a C \ , \text{ y} \quad C^T = -C \ .
\end{aligned} \tag{15}$$

where σ_x, σ_y, and σ_z are Pauli matrices and C stands for the charge conjugation matrix. Using the Jacobi identities we find

$$\chi = \zeta = -1 \quad \mu = \pm\nu \quad \text{y} \quad \tau = \pm 1 \tag{16}$$

The same superalgebra (1-10) is obtained for $D = 3$ if we choose

$$\mu = \nu = 1 \quad \text{and} \quad \tau = +1 \ . \tag{17}$$

In what follows we will use the following superalgebra for $D = 3$

$$[N_{ab}, N_{cd}] = \eta_{ad}N_{bc} + \eta_{bc}N_{ad} - \eta_{ac}N_{bd} - \eta_{bd}N_{ac} \tag{18}$$

$$[L_{ab}, L_{cd}] = \eta_{ad}L_{bc} + \eta_{bc}L_{ad} - \eta_{ac}L_{bd} - \eta_{bd}L_{ac} \tag{19}$$

$$[L_{ab}, L_c] = \eta_{bc}L_a - \eta_{ac}L_b \tag{20}$$

$$[L_a, L_b] = L_{ab} \tag{21}$$

$$[N_{ab}, L_{cd}] = 0 \tag{22}$$

$$[N_{ab}, L_c] = 0. \tag{23}$$

$$\left\{Q'_\alpha, Q'_\beta\right\} = \frac{1}{2}\left[\left(\sigma^{ab}C\right)_{\alpha\beta} L_{ab} - \left(\gamma^a C\right)_{\alpha\beta} L_a\right] \tag{24}$$

$$[L_{ab}, Q'_\alpha] = -\left(\sigma_{ab}Q'\right)_\alpha \tag{25}$$

$$[L_a, Q'_\alpha] = -\frac{1}{2}\left(\gamma_a Q'\right)_\alpha \tag{26}$$

$$[N_{ab}, Q'_\alpha] = 0 \tag{27}$$

where the generators N_{ab} are given by

$$N_{ab} = J_{ab} - \frac{c}{4a^2} Z_{ab}. \tag{28}$$

The set generators $\{N_{ab}\}$ form a basis for the Lorentz algebra $SO(2,1)$. The $\{L_{ab}, L_a\}$ generators, on the other hand, are defined by

$$L_{ab} = \frac{c}{4a^2} Z_{ab},\ L_{a3} = -L_{3a} = \frac{1}{2a} P_a \equiv L_a, \tag{29}$$

which together with the generator $Q'_\alpha \equiv \frac{1}{2a}\sqrt{\frac{c}{2d}} Q_\alpha$ form a basis for the $OSp(2|1) \otimes Sp(2)$ algebra, which is the $N = 1$ AdS superalgebra in $D = 3$ dimensions. Note that the algebra (18-27) is the direct sum $SO(2,1) \oplus OSp(2|1) \otimes Sp(2)$.

III. THE S-EXPANSION OF THE ADS SUPERALGEBRA

In this section we briefly review the general abelian semigroup expansion procedure. Consider a Lie algebra $\mathfrak{g}$ and a finite abelian semigroup $S = \{\lambda_\alpha\}$. According to Theorem 3.1 from Ref. [12], (see also [13], [14], [15], [16]) the direct product $S \times \mathfrak{g}$ is also a Lie algebra. Interestingly, there are cases when it is possible to systematically extract subalgebras from $S \times \mathfrak{g}$. Start by decomposing $\mathfrak{g}$ into a direct sum of subspaces, as in $\mathfrak{g} = \bigoplus_{p\in I} V_p$, where I is a set of indices. The internal structure of $\mathfrak{g}$ can be codified through the subsets $i_{(p,q)} \subset I$ according to $[V_p, V_q] \subset \bigoplus_{r\in i_{(p,q)}} V_r$. When the semigroup S can be decomposed in subsets S_p, $S = \bigcup_{p\in I} S_p$, such that they satisfy the resonant condition $S_p \cdot S_q \subset \bigcap_{r\in i_{(p,q)}} S_r$, then we have that $\mathfrak{G}_{\mathrm{R}} = \bigoplus_{p\in I} S_p \times V_p$ is a 'resonant subalgebra' of $S \times \mathfrak{g}$ (see Theorem 4.2 from Ref. [12]).

An even smaller algebra can be obtained when there is a zero element in the semigroup, i.e., an element $0_S \in S$ such that, for all $\lambda_\alpha \in S$, $0_S \lambda_\alpha = 0_S$. When this is the case, the whole $0_S \times \mathfrak{g}$ sector can be removed from the resonant subalgebra by imposing $0_S \times \mathfrak{g} = 0$. The remaining piece, to which we refer to as 0_S-reduced algebra, continues to be a Lie algebra (see 0_S-reduction and Theorem 6.1 from Ref. [12]).

In order to obtain the $SO(2,1) \oplus OSp(2|1) \otimes Sp(2)$ superalgebra as an S-expansion of anti-de-Sitter $OSp(2|1) \otimes Sp(2)$ superalgebra. The first step consists of splitting the algebra into subspaces, i.e $\mathfrak{g} = OSp(2|1) \otimes Sp(2) = V_0 \oplus V_1 \oplus V_2$, where V_0 corresponds to the Lorentz subalgebra $SO(2,1)$ which is generated by $\widetilde{J}_{ab}$, V_1 corresponds to the supersymmetry translation generated by $\widetilde{Q}_\alpha$ and V_2 corresponds to the AdS boost generated by $\widetilde{P}_a$.

The generators $\widetilde{J}_{ab}$, $\widetilde{Q}_\alpha$, $\widetilde{P}_a$ satisfy the following commutation relation ($D=3$)

$$\left[\tilde{J}_{ab},\tilde{J}_{cd}\right]=\eta_{bc}\tilde{J}_{ad}-\eta_{bd}\tilde{J}_{ac}-\eta_{ac}\tilde{J}_{bd}+\eta_{ad}\tilde{J}_{bc}, \tag{30}$$

$$\left[\tilde{J}_{ab},\tilde{P}_c\right]=\eta_{cb}\tilde{P}_a-\eta_{ca}\tilde{P}_b, \tag{31}$$

$$\left[\tilde{P}_a,\tilde{P}_b\right]=\tilde{J}_{ab}, \tag{32}$$

$$\left[\tilde{J}_{ab},\tilde{Q}\right]=-\sigma_{ab}\tilde{Q}, \tag{33}$$

$$\left[\tilde{P}_a,\tilde{Q}\right]=-\frac{1}{2}\gamma_a\tilde{Q}, \tag{34}$$

$$\left\{\tilde{Q}_\alpha,\tilde{Q}_\beta\right\}=\frac{1}{2}\left[\left(\sigma^{ab}C\right)_{\alpha\beta}\tilde{J}_{ab}-\left(\gamma^aC\right)_{\alpha\beta}\tilde{P}_a\right]. \tag{35}$$

which have the following subspace structure

$$\begin{aligned}&[V_0,V_0]\subset V_0,\quad [V_0,V_2]\subset V_2,\ \ [V_2,V_2]\subset V_0,\\ &[V_0,V_1]\subset V_1,\quad [V_2,V_1]\subset V_1,\ \ [V_1,V_1]\subset V_0\oplus V_2.\end{aligned} \tag{36}$$

The second step consists of finding an abelian semigroup S which can be partitioned in a "resonant" way with respect to (36). We shall consider the expansion using two different semigroups (see [2]).

A. Semigroup S_{S3}

Let us consider first the semigroup $S_{S3}=\left\{\overline{\lambda}_0,\overline{\lambda}_1,\overline{\lambda}_2,\overline{\lambda}_3\right\}$ defined by the product table:

	$\overline{\lambda}_0$	$\overline{\lambda}_1$	$\overline{\lambda}_2$	$\overline{\lambda}_3$
$\overline{\lambda}_0$	$\overline{\lambda}_2$	$\overline{\lambda}_3$	$\overline{\lambda}_0$	$\overline{\lambda}_3$
$\overline{\lambda}_1$	$\overline{\lambda}_3$	$\overline{\lambda}_1$	$\overline{\lambda}_3$	$\overline{\lambda}_3$
$\overline{\lambda}_2$	$\overline{\lambda}_0$	$\overline{\lambda}_3$	$\overline{\lambda}_2$	$\overline{\lambda}_3$
$\overline{\lambda}_3$	$\overline{\lambda}_3$	$\overline{\lambda}_3$	$\overline{\lambda}_3$	$\overline{\lambda}_3$

(37)

A straightforward but important observation is that, for each $\lambda_\alpha\in S$, $\overline{\lambda}_3\overline{\lambda}_\alpha=\overline{\lambda}_3$, so that $\overline{\lambda}_3$ plays the role of the zero element inside S_{S3}. Consider now the partition $S=S_0\cup S_1\cup S_2$, with

$$S_0=\left\{\overline{\lambda}_1,\overline{\lambda}_2,\overline{\lambda}_3\right\}, \tag{38}$$

$$S_1=\left\{\overline{\lambda}_0,\overline{\lambda}_3\right\}, \tag{39}$$

$$S_2=\left\{\overline{\lambda}_2,\overline{\lambda}_3\right\}. \tag{40}$$

This partition is said to be resonant, since it satisfies [compare eqs. (36) with eqs. (41)]

$$\begin{aligned} &S_0 \cdot S_0 \subset S_0, \quad S_0 \cdot S_2 \subset S_2, \ S_2 \cdot S_2 \subset S_0 \\ &S_0 \cdot S_1 \subset S_1, \quad S_2 \cdot S_1 \subset S_1, \ S_1 \cdot S_1 \subset S_0 \cap S_2 \end{aligned} \tag{41}$$

Using Theorem 4.2 from Ref. [12] one sees that the algebra

$$\mathfrak{G}_{\rm R} = W_0 \oplus W_1 \oplus W_2, \tag{42}$$

is a *resonant subalgebra* of $S_{\rm S3} \times \mathfrak{g}$, where

$$W_0 = (S_0 \times V_0) = \left\{\overline{\lambda}_1, \overline{\lambda}_2, \overline{\lambda}_3\right\} \otimes \{J_{ab}\} = \left\{\overline{\lambda}_1 \tilde{J}_{ab}, \overline{\lambda}_2 \tilde{J}_{ab}, \overline{\lambda}_3 \tilde{J}_{ab}\right\} \tag{43}$$

$$W_1 = (S_1 \times V_1) = \left\{\overline{\lambda}_0, \overline{\lambda}_3\right\} \otimes \left\{\widetilde{Q}_\alpha\right\} = \left\{\overline{\lambda}_0 \widetilde{Q}_\alpha, \overline{\lambda}_3 \widetilde{Q}_\alpha\right\} \tag{44}$$

$$W_2 = (S_2 \times V_2) = \left\{\overline{\lambda}_2, \overline{\lambda}_3\right\} \otimes \left\{\tilde{P}_a\right\} = \left\{\overline{\lambda}_2 \tilde{P}_a, \overline{\lambda}_3 \tilde{P}_a\right\} \tag{45}$$

Finally, we impose the condition $\lambda_3 \times \mathfrak{g} = 0$ on $\mathfrak{G}_{\rm R}$ and relabel its generators in the form $J_{ab,1} = \overline{\lambda}_1 \tilde{J}_{ab}$; $J_{ab,2} = \overline{\lambda}_2 \tilde{J}_{ab}$; $Q_{\alpha,0} = \overline{\lambda}_0 \widetilde{Q}_\alpha$; $P_{a,2} = \overline{\lambda}_2 \tilde{P}_a$. This procedure leads us to the following commutation relations:

$$\begin{aligned} [J_{ab,1}, J_{cd,1}] &= \overline{\lambda}_1 \overline{\lambda}_1 \left[J_{ab}, J_{cd}\right] = \overline{\lambda}_1 \left[\tilde{J}_{ab}, \tilde{J}_{cd}\right] \\ &= \eta_{ad} J_{bc,1} + \eta_{bc} J_{ad,1} - \eta_{ac} J_{bd,1} - \eta_{bd} J_{ac,1} \end{aligned} \tag{46}$$

$$\begin{aligned} [J_{ab,2}, J_{cd,2}] &= \overline{\lambda}_2 \overline{\lambda}_2 \left[\tilde{J}_{ab}, \tilde{J}_{cd}\right] = \overline{\lambda}_2 \left[\tilde{J}_{ab}, \tilde{J}_{cd}\right] \\ &\eta_{ad} J_{bc,2} + \eta_{bc} J_{ad,2} - \eta_{ac} J_{bd,2} - \eta_{bd} J_{ac,2} \end{aligned} \tag{47}$$

$$[J_{ab,1}, J_{cd,2}] = \overline{\lambda}_2 \overline{\lambda}_2 \left[\tilde{J}_{ab}, \tilde{J}_{cd}\right] = \overline{\lambda}_3 \left[\tilde{J}_{ab}, \tilde{J}_{cd}\right] = 0 \tag{48}$$

$$[J_{ab,1}, P_{c,2}] = \overline{\lambda}_1 \overline{\lambda}_2 \left[\tilde{J}_{ab}, P_c\right] = \overline{\lambda}_3 \left[\tilde{J}_{ab}, P_c\right] = 0 \tag{49}$$

$$\begin{aligned} [J_{ab,2}, P_{c,2}] &= \lambda_2 \lambda_2 \left[\tilde{J}_{ab}, \tilde{P}_c\right] = \lambda_2 \left[\tilde{J}_{ab}, \tilde{P}_c\right] \\ &= \eta_{bc} P_{a,2} - \eta_{ac} P_{b,2} \end{aligned} \tag{50}$$

$$[P_{a,2}, P_{b,2}] = \overline{\lambda}_2 \overline{\lambda}_2 \left[\tilde{P}_a, \tilde{P}_b\right] = \overline{\lambda}_2 \left[\tilde{P}_a, \tilde{P}_b\right] = \overline{\lambda}_2 \tilde{J}_{ab} = J_{ab,2} \tag{51}$$

$$[J_{ab,1}, Q_{\alpha,0}] = \overline{\lambda}_1 \overline{\lambda}_0 \left[\tilde{J}_{ab}, \widetilde{Q}_\alpha\right] = \overline{\lambda}_3 \left[\tilde{J}_{ab}, \widetilde{Q}_\alpha\right] = 0 \tag{52}$$

$$[J_{ab,2}, Q_{\alpha,0}] = \overline{\lambda}_2\overline{\lambda}_0 \left[\boldsymbol{\tilde{J}}_{ab}, \widetilde{Q}_\alpha\right] = \overline{\lambda}_0 \left[\tilde{J}_{ab}, \widetilde{Q}_\alpha\right] \\ = -\left(\sigma_{ab}\widetilde{Q}_0\right)_\alpha \tag{53}$$

$$[P_{a,2}, Q_{\alpha,0}] = \overline{\lambda}_2\overline{\lambda}_0 \left[\tilde{P}_a, \widetilde{Q}_\alpha\right] = \overline{\lambda}_0 \left[\tilde{P}_a, \widetilde{Q}_\alpha\right] \\ = -\frac{1}{2}\left(\gamma_a\widetilde{Q}_0\right)_\alpha \tag{54}$$

$$\{Q_{\alpha,0}, Q_{\beta,0}\} = \overline{\lambda}_0\overline{\lambda}_0 \left\{\widetilde{Q}_\alpha, \widetilde{Q}_\beta\right\} = \overline{\lambda}_2 \left\{\widetilde{Q}_\alpha, \widetilde{Q}_\beta\right\} \\ = \frac{1}{2}\left[\left(\sigma^{ab}C\right)_{\alpha\beta} J_{ab,2} - \left(\gamma^a C\right)_{\alpha\beta} P_{a,2}\right] \tag{55}$$

where we have used the commutation relations of the AdS superalgebra and the law of multiplication of the semigroup (37).

The identification $N_{ab} = J_{ab,1}$; $L_{ab} = J_{ab,2}$; $L_a = P_{a,2}$; $Q'_\alpha = Q_{\alpha,0}$; shows that the superalgebra (46-55), obtained by expansion of the AdS superalgebra $OSp\left(2|1\right) \otimes Sp\left(2\right)$, is the $SO\left(2,1\right) \oplus OSp\left(2|1\right) \otimes Sp\left(2\right)$ AdS–Lorentz superalgebra (18-27) obtained by semisimple extension of the Poincare superalgebra in ref. ([1], [5]).

B. Semigroup S_{S2}

Let us consider the semigroup $S_{S2} = \{\lambda_0, \lambda_1, \lambda_2\}$ defined by the multiplication law (see [2])

$$\lambda_\alpha\lambda_\beta = \begin{cases} \lambda_{\alpha+\beta}, \text{ if } \alpha+\beta \leq 2 \\ \lambda_{\alpha+\beta-2}, \text{ if } \alpha+\beta > 2 \end{cases} \tag{56}$$

Take now the partition $S = S_0 \cup S_1 \cup S_2$, with

$$S_0 = \{\lambda_0, \lambda_2\}, \tag{57}$$

$$S_1 = \{\lambda_1\}, \tag{58}$$

$$S_2 = \{\lambda_2\}. \tag{59}$$

This partition is said to be resonant, since it satisfies

$$S_0 \cdot S_0 \subset S_0, \tag{60}$$

$$S_0 \cdot S_1 \subset S_1, \tag{61}$$

$$S_0 \cdot S_2 \subset S_2, \tag{62}$$

$$S_1 \cdot S_1 \subset S_0 \cap S_2, \tag{63}$$

$$S_1 \cdot S_2 \subset S_1, \tag{64}$$

$$S_2 \cdot S_2 \subset S_2. \tag{65}$$

As in the previous case, Theorem 4.2 from Ref. [12] now assures us that

$$\mathfrak{G}_{\mathrm{R}} = W_0 \oplus W_1 \oplus W_2, \tag{66}$$

is a *resonant subalgebra* of $S_{\mathrm{S2}} \times \mathfrak{g}$, where

$$W_0 = (S_0 \times V_0) = \{\lambda_0, \lambda_2\} \otimes \left\{\tilde{J}_{ab}\right\} = \left\{\lambda_0 \tilde{J}_{ab}, \lambda_2 \tilde{J}_{ab}\right\} \tag{67}$$

$$W_1 = (S_1 \times V_1) = \{\lambda_1\} \otimes \left\{\widetilde{Q}_\alpha\right\} = \left\{\lambda_1 \widetilde{Q}_\alpha\right\} \tag{68}$$

$$W_2 = (S_2 \times V_2) = \{\lambda_2\} \otimes \left\{\tilde{P}_a\right\} = \left\{\lambda_2 \tilde{P}_a\right\} \tag{69}$$

Relabelling the generators of the resonant sub-superalgebra as

$$J_{ab} = \lambda_0 \tilde{J}_{ab}, \tag{70}$$

$$Q = \lambda_1 \tilde{Q}, \tag{71}$$

$$Z_{ab} = \lambda_2 \tilde{J}_{ab}, \tag{72}$$

$$P_a = \lambda_2 \tilde{P}_a. \tag{73}$$

we are left with the following commutation relations:

$$[J_{ab}, J_{cd}] = \eta_{bc} J_{ad} - \eta_{bd} J_{ac} - \eta_{ac} J_{bd} + \eta_{ad} J_{bc}, \tag{74}$$

$$[J_{ab}, Z_{cd}] = \eta_{bc} Z_{ad} - \eta_{bd} Z_{ac} - \eta_{ac} Z_{bd} + \eta_{ad} Z_{bc}, \tag{75}$$

$$[Z_{ab}, Z_{cd}] = \eta_{bc} Z_{ad} - \eta_{bd} Z_{ac} - \eta_{ac} Z_{bd} + \eta_{ad} Z_{bc}, \tag{76}$$

$$[J_{ab}, P_c] = \eta_{cb} P_a - \eta_{ca} P_b, \tag{77}$$

$$[Z_{ab}, P_c] = \eta_{cb} P_a - \eta_{ca} P_b, \tag{78}$$

$$[P_a, P_b] = Z_{ab}, \tag{79}$$

$$[J_{ab}, Q] = -\sigma_{ab} Q, \tag{80}$$

$$[Z_{ab}, Q] = -\sigma_{ab} Q, \tag{81}$$

$$[P_a, Q] = -\frac{1}{2} \gamma_a Q, \tag{82}$$

$$\{Q_\alpha, Q_\beta\} = \frac{1}{2} \left[\left(\sigma^{ab} C\right)_{\alpha\beta} Z_{ab} - \left(\gamma^a C\right)_{\alpha\beta} P_a \right]. \tag{83}$$

which is the supersymmetric extension of the so called AdS–Lorentz algebra.

The relationship between multiplication tables of semigroups S_{S3} and S_{S2} is given by (see [2])

$$\bar{\lambda}_1 = \lambda_0 - \lambda_2, \tag{84}$$

$$\bar{\lambda}_2 = \lambda_2, \tag{85}$$

$$\bar{\lambda}_0 = \lambda_1, \tag{86}$$

This amounts to a change of basis in S_{S2} and leads to the following multiplication table (see [2]):

	$\bar{\lambda}_0$	$\bar{\lambda}_1$	$\bar{\lambda}_2$
$\bar{\lambda}_0$	$\bar{\lambda}_2$	0	$\bar{\lambda}_0$
$\bar{\lambda}_1$	0	$\bar{\lambda}_1$	0
$\bar{\lambda}_2$	$\bar{\lambda}_0$	0	$\bar{\lambda}_2$

(87)

The Maxwell superalgebra can be obtained from the AdS–Lorentz superalgebra (74)-(83) by means of a generalized Inönü–Wigner contraction [17]. In fact, the rescaling

$$Z_{ab} \to \lambda^2 Z_{ab} \quad , \qquad P_a \to \lambda P_a \quad , \text{ y} \qquad Q_\alpha \to \lambda Q_\alpha \tag{88}$$

in (74)-(83) provides, in the limit $\lambda \to \infty$, the $N = 1$ Maxwell superalgebra (see [17])

$$[J_{ab}, J_{cd}] = \eta_{bc} J_{ad} - \eta_{bd} J_{ac} - \eta_{ac} J_{bd} + \eta_{ad} J_{bc}, \tag{89}$$

$$[J_{ab}, P_c] = \eta_{cb} P_a - \eta_{ca} P_b, \tag{90}$$

$$[P_a, P_b] = Z_{ab}, \tag{91}$$

$$[Z_{ab}, Z_{cd}] = 0, \tag{92}$$

$$[J_{ab}, Z_{cd}] = \eta_{bc} Z_{ad} - \eta_{bd} Z_{ac} - \eta_{ac} Z_{bd} + \eta_{ad} Z_{bc}, \tag{93}$$

$$[Z_{ab}, P_c] = 0, \tag{94}$$

$$[J_{ab}, Q_\alpha] = -(\sigma_{ab} Q)_\alpha, \tag{95}$$

$$[Z_{ab}, Q_\alpha] = 0, \tag{96}$$

$$[P_a, Q_\alpha] = 0, \tag{97}$$

$$\{Q_\alpha, Q_\beta\} = \frac{1}{2} \left(\sigma^{ab} C\right)_{\alpha\beta} Z_{ab}. \tag{98}$$

C. AdS–Lorentz Casimir operators

In ref. [18] was found that the invariant tensors for the AdS superalgebra in $(2 + 1)$ dimensions are given by

$$\langle J_{ab} J_{cd} \rangle = \tilde{\mu}_0 \left(\eta_{ad}\eta_{bc} - \eta_{ac}\eta_{bd}\right) \tag{99}$$

$$\langle J_{ab} P_c \rangle = \tilde{\mu}_1 \epsilon_{abc} \tag{100}$$

$$\langle P_a P_b \rangle = \tilde{\mu}_0 \eta_{ab} \tag{101}$$

$$\langle Q_\alpha Q_\beta \rangle = (\tilde{\mu}_0 - \tilde{\mu}_1) C_{\alpha\beta}. \tag{102}$$

which are also the components of the Killing metric for the AdS superalgebra

$$k_{ab,cd} = \tilde{\mu}_0 \left(\eta_{ad}\eta_{bc} - \eta_{ac}\eta_{bd}\right) \tag{103}$$

$$k_{ab,c} = k_{c,ab} = \tilde{\mu}_1 \epsilon_{abc} \tag{104}$$

$$k_{a,b} = \tilde{\mu}_0 \eta_{ab} \tag{105}$$

$$k_{\alpha,\beta} = (\tilde{\mu}_0 - \tilde{\mu}_1) C_{\alpha\beta} \tag{106}$$

Following the procedure developed in ref. [2], we have

$$T_A = k_{AC} k^{CB} T_B \tag{107}$$

i.e.,

$$J_{ab} = k_{ab,mn}k^{mn,cd}J_{cd} + k_{ab,m}k^{m,cd}J_{cd} + k_{ab,m}k^{m,c}P_c \tag{108}$$

so that, $k^{a,b} = 0$ and

$$J_{ab} = 2\tilde{\mu}_0 k_{ba}{}^{cd}J_{cd} + \tilde{\mu}_1\epsilon_{abm}k^{m,cd}J_{cd} \tag{109}$$

If

$$k^{ab,cd} = \alpha\eta^{[ab][cd]} \tag{110}$$

$$k^{m,cd} = \beta\epsilon^{mcd} \tag{111}$$

then

$$J_{ab} = 2\left(2\alpha\tilde{\mu}_0 - \beta\tilde{\mu}_1\right)J_{ab} \tag{112}$$

so that

$$\beta = \frac{1}{\tilde{\mu}_1}\left(2\alpha\tilde{\mu}_0 - \frac{1}{2}\right) \tag{113}$$

In the same way we obtain

$$P_a = k_{a,m}k^{m,b}P_b + k_{a,mn}k^{mn,b}P_b + k_{a,mn}k^{mn,cd}J_{cd} \tag{114}$$

so that, $\alpha = 0$. Since $k^{a,b} = 0$, we have

$$P_a = \tilde{\mu}_1\epsilon_{amn}k^{mn,b}P_b \tag{115}$$

Since

$$Q_\alpha = k_{\alpha,\gamma}k^{\gamma,\beta}Q_\beta \tag{116}$$

$$= (\tilde{\mu}_0 - \tilde{\mu}_1)C_{\alpha\gamma}k^{\gamma,\beta}Q_\beta \tag{117}$$

we have,

$$k^{\alpha,\beta} = -\frac{1}{\tilde{\mu}_0 - \tilde{\mu}_1}C^{\alpha\beta} \tag{118}$$

Summarizing we have found that

$$k^{ab,c} = -\frac{1}{2\tilde{\mu}_1}\epsilon^{abc} \tag{119}$$

$$k^{\alpha,\beta} = -\frac{1}{\tilde{\mu}_0 - \tilde{\mu}_1}C^{\alpha\beta} \tag{120}$$

This means that the Casimir Operator for the AdS superalgebra in $d = 3$ is given by

$$C = -\frac{1}{2\tilde{\mu}_1}\epsilon^{abc}\left(J_{ab}P_c + P_cJ_{ab}\right) - \frac{1}{\tilde{\mu}_0 - \tilde{\mu}_1}C^{\alpha\beta}Q_\alpha Q_\beta \tag{121}$$

$$= -\frac{1}{2\tilde{\mu}_1}\epsilon^{abc}\left(J_{ab}P_c + P_cJ_{ab}\right) - \frac{1}{\tilde{\mu}_0 - \tilde{\mu}_1}Q\bar{Q} \tag{122}$$

From [2] we know that the Casimir operator for S-expanded Lie algebras is given by

$$C = m^{\alpha\beta}C^{AB}T_{(A,\alpha)}T_{(B,\beta)}, \tag{123}$$

where C^{AB} are the components of the Casimir operator for the original algebra $\mathfrak{g}$ and $m^{\alpha\beta}$ is the inverse of the matrix $m_{\alpha\beta} = \alpha_\gamma K_{\alpha\beta}{}^{\gamma}$. The two-selectors $K_{\alpha\beta}{}^{\gamma}$ of S_{S2} can be represented as (see [2])

$$K_{\alpha\beta}{}^{0} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},\ K_{\alpha\beta}{}^{1} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix},\ K_{\alpha\beta}{}^{2} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}. \tag{124}$$

This implies that the metric $m_{\alpha\beta}$ for S_{S2} reads

$$m_{\alpha\beta} = \alpha_\lambda K_{\alpha\beta}{}^{\lambda} = \begin{pmatrix} \alpha_0 & \alpha_1 & \alpha_2 \\ \alpha_1 & \alpha_2 & \alpha_1 \\ \alpha_2 & \alpha_1 & \alpha_2 \end{pmatrix} \tag{125}$$

The inverse matrix is then given by

$$m^{\alpha\beta} = \frac{1}{\det m_{\alpha\beta}} \begin{pmatrix} \alpha_2^2 - \alpha_1^2 & 0 & -\left(\alpha_2^2 - \alpha_1^2\right) \\ 0 & \alpha_2\left(\alpha_0 - \alpha_2\right) & -\alpha_1\left(\alpha_0 - \alpha_2\right) \\ -\left(\alpha_2^2 - \alpha_1^2\right) & -\alpha_1\left(\alpha_0 - \alpha_2\right) & \alpha_0\alpha_2 - \alpha_1^2 \end{pmatrix}, \tag{126}$$

where

$$\det m_{\alpha\beta} = \left(\alpha_0 - \alpha_2\right)\left(\alpha_2 + \alpha_1\right)\left(\alpha_2 - \alpha_1\right) \neq 0. \tag{127}$$

so that the quadratic Casimir Operator for the semi-simple extended Poincaré superalgebra in $d = 3$ have the form

$$\begin{aligned} C = &\left(\alpha_2^2 - \alpha_1^2\right)\frac{1}{2\tilde{\mu}_1}\epsilon^{abc}\left(J_{ab}P_c + P_cJ_{ab}\right) \\ &- \left(\alpha_0\alpha_2 - \alpha_1^2\right)\frac{1}{2\tilde{\mu}_1}\epsilon^{abc}\left(Z_{ab}P_c + P_cZ_{ab}\right) \\ &- \left(\alpha_2\alpha_0 - \alpha_2^2\right)\frac{1}{\tilde{\mu}_0 - \tilde{\mu}_1}Q\bar{Q} \end{aligned} \tag{128}$$

Defining

$$\alpha = -\left(\alpha_2\alpha_0 - \alpha_2^2\right) \tag{129}$$

$$\beta = -\left(\alpha_2\alpha_0 - \alpha_1^2\right) \tag{130}$$

we obtain

$$\begin{aligned} C = &\frac{\alpha}{\tilde{\mu}_1}\left[\frac{1}{2}\epsilon^{abc}\left(J_{ab}P_c + P_cJ_{ab}\right) + \frac{1}{1-\frac{\tilde{\mu}_0}{\tilde{\mu}_1}}Q\bar{Q}\right] \\ &+ \frac{\beta}{2\tilde{\mu}_1}\epsilon^{abc}\left[Z_{ab}P_c + P_cZ_{ab} - J_{ab}P_c - P_cJ_{ab}\right] \end{aligned} \tag{131}$$

This means that the semi-simple extended Poincaré superalgebra in $d = 3$ posses two independent Casimir operators

$$C_1 = \frac{1}{2}\epsilon^{abc}\left(J_{ab}P_c + P_cJ_{ab}\right) + \frac{1}{1-\frac{\tilde{\mu}_0}{\tilde{\mu}_1}}Q\bar{Q}, \tag{132}$$

$$C_2 = \epsilon^{abc}\left[Z_{ab}P_c + P_cZ_{ab} - J_{ab}P_c - P_cJ_{ab}\right]. \tag{133}$$

IV. ADS–LORENTZ CHERN-SIMONS SUPERGRAVITY IN $D = 3$

In this section we present the construction of the Chern-Simons supergravity action in $(2+1)-$dimensions.

A. The Invariant Tensor

In Ref. [18] was found that the invariant tensors for the AdS superalgebra in $(2+1)-$dimensions are given by eqs. (99-102). Now, following Theorem 7.2 from Ref. [12] ensures us that the only nonzero components of a symmetric invariant tensor for the AdS–Lorentz superalgebra are:

$$\langle N_{ab}N_{cd}\rangle = \alpha_0\left(\eta_{ad}\eta_{bc} - \eta_{ac}\eta_{bd}\right) \tag{134}$$

$$\langle L_{ab}L_{cd}\rangle = \alpha_2\left(\eta_{ad}\eta_{bc} - \eta_{ac}\eta_{bd}\right) \tag{135}$$

$$\langle L_{ab}L_{c3}\rangle = \alpha_1\epsilon_{abc} \tag{136}$$

$$\langle L_{a3}L_{b3}\rangle = \alpha_2\eta_{ab} \tag{137}$$

$$\left\langle Q'_\alpha Q'_\beta\right\rangle = (\alpha_2 - \alpha_1)C_{\alpha\beta}. \tag{138}$$

B. AdS–Lorentz invariant supergravity action

Let us now turn to gauging the AdS–Lorentz superalgebra. To this end we write down gauge field A, valued in this superalgebra

$$A = \frac{1}{2}\varpi^{ab}N_{ab} + \frac{1}{2}\omega^{ab}L_{ab} + \omega^{a3}L_{a3} + \psi^{\alpha}Q'_{\alpha}. \tag{139}$$

The Chern–Simons Lagrangian in $(2+1)-$dimensions [19], [21], is given by

$$L_{CS}^{2+1} = 2k\int_0^1 dt\left\langle A\left(tdA + t^2A^2\right)\right\rangle = k\left\langle A\left(dA + \frac{2}{3}A^2\right)\right\rangle. \tag{140}$$

Inserting (139) into (140) and using (134-138), the Chern–Simons action for the AdS–Lorentz superalgebra, in the basis $\{N_{ab}, L_{ab}, L_{a3}, \psi^{\alpha}\}$, is given by

$$\begin{aligned} S_{CS}^{(2+1)} =& k\int \frac{1}{2}\alpha_0\varpi^a{}_c\left(d\varpi^c{}_a + \frac{2}{3}\varpi^c{}_d\varpi^d{}_a\right) && (141)\\ &+ \alpha_1\epsilon_{abc}\left(R^{ab}\omega^{c3} + \frac{1}{3}\omega^{a3}\omega^{b3}\omega^{c3}\right) + (\alpha_1-\alpha_2)\psi^{\alpha}D\psi_{\alpha} && (142)\\ &- \alpha_2 D\omega^{a3}\omega^3{}_a + \frac{1}{2}\alpha_2\omega^a{}_c\left(d\omega^c{}_a + \frac{2}{3}\omega^c{}_d\omega^d{}_a\right) - d\left\{\alpha_1\frac{1}{2}\varepsilon_{abc}\omega^{ab}\omega^{c3}\right\}. && (143)\end{aligned}$$

Here

$$R^{ab} = d\omega^{ab} + \omega^a{}_c\omega^{cb} \tag{144}$$

$$D\psi = d\psi + \frac{1}{4}\omega^{ab}\gamma_{ab}\psi \tag{145}$$

$$D\omega^{a3} = d\omega^{a3} + \omega^a{}_b\omega^{b3} \tag{146}$$

Identifying $\omega^{a3} = e^a/l$, where e^a is the vielbein and l is a parameter with dimensions of length, we find

$$\begin{aligned} S_{CS}^{(2+1)} =& k\int_M \frac{1}{2}\alpha_0\varpi^a{}_c\left(d\varpi^c{}_a + \frac{2}{3}\varpi^c{}_d\varpi^d{}_a\right) && (147)\\ &+ \frac{\alpha_1 k}{l}\left[\int_M \epsilon_{abc}\left(R^{ab}e^c + \frac{1}{3l^2}e^ae^be^c\right) + \bar{\psi}^{\alpha}D\psi_{\alpha} - \frac{1}{2}\int_{\partial M}\varepsilon_{abc}\omega^{ab}e^c\right] && (148)\\ &+ \frac{1}{2}\alpha_2 k\int_M\left[\omega^a{}_c\left(d\omega^c{}_a + \frac{2}{3}\omega^c{}_d\omega^d{}_a\right) + \frac{2}{l^2}e_aT^a - 2\bar{\psi}^{\alpha}D\psi_{\alpha}\right]. && (149)\end{aligned}$$

where we have used $D_{\omega}\omega^{a3} = (De^a)/l = T^a/l$.

V. COMMENTS

We have shown that: (i) The AdS–Lorentz superalgebra $SO(2,1) \oplus OSp(2|1) \otimes Sp(2)$ of Ref. [1] can be obtained from the AdS–superalgebra $OSp(2|1) \otimes Sp(2)$ via the $S-$expansion procedure [12] with an appropriate semigroup. (ii) The above mentioned expansion method allows to obtain the Casimir operators and the invariant tensor for the AdS–Lorentz superalgebra, which allows to construct an action for Chern–Simons supergravity in $(2+1)-$dimensions, compare with the action of ref. [20]. The resulting theory contain the so called exotic Lagrangian [21] and the minimal supersymmetric extension of the topological Mielke-Baekler action for three-dimensional gravity [22], [23].

This work was supported in part by FONDECYT through Grants # 1130653. Two of the authors (O.F and O.V) were supported by grants from the Comisión Nacional de Investigación Científica y Tecnológica CONICYT and from the Universidad de Concepción, Chile.

---

[1] D. V. Soroka and V. A. Soroka, Adv. High Energy Phys. 2009 (2009) 234147. ArXiv:hep-th/0605251.

[2] J. Díaz, O. Fierro, F. Izaurieta, N. Merino, E. Rodríguez, P. Salgado, O. Valdivia, J. Phys. A: Math. Theor. 45 (2012) 255207.

[3] H. Bacry, P. Combe and J. L. Richard, Nuovo Cim. A67 (1970) 267-299.

[4] R. Schrader, Fortsch. Phys. 20 (1972) 701-734.

[5] D. V. Soroka and V. A. Soroka, Phys. Lett. B607 (2005) 302; hep-th/0410012.

[6] S. Bonanos, J. Gomis, K. Kamimura and J. Lukierski, Phys. Rev. Lett. 104 (2010) 090401. Arxiv hep-th/0911.5072

[7] K. Kamimura and J. Lukierski, Phys. Lett. B707 (2012) 292-297.

[8] S. Bonanos, J. Gomis, K. Kamimura and J. Lukierski, J. Math. Phys. 51 (2010) 102301. ArXiv:hep-th/1005.3714.

[9] J. A. de Azcárraga, K. Kamimura, and J. Lukierski, Phys.Rev. D83 (2011) 124036. ArXiv hep-th/1012.4402

[10] R. Durka, J. Kowalski-Glikman and M. Szczachor, Mod. Phys. Lett. A26 (2011) 2689.

ArXiv:hep-th/1107.4728

[11] R. Durka, J. Kowalski-Glikman and M. Szczachor, Mod. Phys. Lett. A27 (2012) 1250023.

[12] F. Izaurieta, E. Rodriguez, P. Salgado, J. Math. Phys.47. (2006) 123512. ArXiv:hep-th/0606215.

[13] M. Hatsuda, M. Sakaguchi, Prog. Theor. Phys. **109** (2003) 853. arXiv: hep-th/0106114.

[14] J. A. de Azcárraga, J. M. Izquierdo, M. Picón, O. Varela, Nucl. Phys. B **662** (2003) 185. arXiv: hep-th/0212347.

[15] J. A. de Azcárraga, J. M. Izquierdo, M. Picón, O. Varela, Class. Quant. Grav. **21** (2004) S1375. arXiv: hep-th/0401033.

[16] J. A. de Azcárraga, J. M. Izquierdo, M. Picón, O. Varela, Int. J. Theor. Phys. **46** (2007) 2738. arXiv: hep-th/0703017.

[17] J. Lukierski, Proc. Steklov Inst. Math. 272 (2011) 183. arXiv 1007.3405

[18] R. Troncoso, "Supergravedad en Dimensiones Impares" Doctoral Thesis, Universidad de Chile, Santiago (1996).

[19] A.H. Chamseddine, Phys. Lett. B 233 (1989) 291; Nucl. Phys. B 346 (1990) 213.

[20] A. Achucarro and P. K. Townsend, A Chern-Simons Action For Three- Dimensional Anti-De Sitter Supergravity Theories, Phys. Lett. B 180, 89 (1986).

[21] J. Zanelli, Lectures on Chern-Simons (super)gravities. Second Edition (February 2008). ArXiv:hep-th/0502193

[22] E.W. Mielke and P. Baekler, Phys. Lett. A 156 (1991) 399

[23] A. Giacomini, R. Troncoso and S. Willison, Three-dimensional supergravity reloaded, Class.Quant.Grav. 24 (2007) 2845-2860. Arxiv:hep-th/0610077.